

Original Research

KNOWLEDGE, ATTITUDE AND PRACTICE OF KANGAROO MOTHER CARE AMONG POSTNATAL MOTHERS IN A TERTIARY CARE CENTER OF HARARI REGION, HARAR, ETHIOPIA

Arif H. Jamie

Harar Health Science College, Department of Pediatrics Nursing, East Ethiopia, Ethiopia

Received: 27 October 2020 | Accepted: 11 November 2020

DOI: <https://dx.doi.org/10.36685/phi.v6i4.355>

Correspondence:

Arif Hussien, MSc

Harar Health Science College, Department of Pediatrics Nursing, East Ethiopia, Ethiopia

PoBox 806, Tel.0921843442

Email: arifhussen.ah@gmail.com

Copyright: © 2020 the author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Kangaroo Mother Care (KMC) is a method was invented by Dr. Rey in 1978 and developed by Dr. Martinez and Dr. Navarrete until 1994. It is holding a small nappy neonate in skin to skin contact, prone and upright on the maternal chest. At first it was developed to alleviate overcrowding, and insufficient resources in neonatal intensive care units. But today it is formally approved by WHO.

Objective: The aim of this study was to assess knowledge, attitude and practice of Kangaroo mother care among postnatal mothers in a tertiary care center of Harari region, Harar, Ethiopia, 2020.

Method: An institution based cross-sectional study method was conducted in Hiwot-Fana Specialized Referral Hospital in Harar town. Sample size was calculated by using single population proportion sample formula and the final sample size was 166. The study subject was selected by using consecutive sampling method, and adopted data collection tool was used. Data were analyzed using SPSS version 20. Descriptive analyses were performed and bivariate analyses were used to find out the association of independent variables.

Results: There was 82.53% of mothers had good knowledge, 82.53% had positive attitude towards KMC, 32.12% practiced KMC correctly. The participants also reported the benefits of KMC, such as protection from infection, warmth to the baby, exclusive breast feeding, weight gain and growth, early discharge from hospital, safety and love.

Conclusion: The finding of this study showed that there was a clear gap in knowledge and practice. Therefore, it is very important to have health education sessions during ANC follow-up on KMC. A qualitative research to explore temporal relationship is also recommended.

Keywords: knowledge, attitude, practice, Kangaroo mother care, Harar, Ethiopia

BACKGROUND

Kangaroo Mother Care (KMC) is a method was invented by Dr. Rey in 1978 and developed by Dr. Martinez and Dr. Navarrete until 1994. It is holding a small nappy neonate in skin to skin contact, prone and upright on the maternal chest. At first it was developed to alleviate overcrowding, and insufficient resources in

neonatal intensive care units. But today it is formally approved by World Health Organization ([WHO, 2003](#)).

The key components of kangaroo mother care are skin-to-skin contact of the infant on the mother's chest, optimal breast feeding and support to the mother and infant. It has more advantages over usual new born care for mother and infant, such

as effective body temperature control, exclusive breastfeeding, preventing infection, mother-infant bonding, early discharge, decreased mortality and morbidity. It is also an answer to the separation of mothers and infants ([Conde-Agudelo & Díaz-Rossello, 2016](#); [M Ludington-Hoe, 2011](#); [Nygqvist et al., 2010](#)).

Globally, each year nearly 7.7 million children under five years die; among this, about 3.1 million of the newborns die during the neonatal period and almost all (99%) deaths occur in the developing countries ([Rajaratnam et al., 2010](#)). World Health Organization estimation, neonatal deaths account for 45% of the under-five deaths ([WHO, 2017](#)). Three-quarter of the neonatal deaths takes place in the first seven days of birth and one-third of deaths take place in the first 24 h of birth ([Aker, Dawson, & Sibbritt, 2016](#); [Lawn, Cousens, Zupan, & Team](#)). The three leading causes of neonatal mortality and morbidity worldwide are infections (35%), preterm birth and LBW (28%) and asphyxia (23%) ([WHO & UNICEF, 2009](#)).

Globally, babies with low birth weight accounts 25 million and preterm accounts 15 million nearly all of them (96%) are in developing countries. Africa accounts the world's population of 12% and 25% of the world's new-born deaths takes place there ([WHO, 2003](#); [WHO & UNICEF, 2004](#)).

Neonatal hypothermia is an important confront associated with morbidity and mortality ([Trevisanuto et al., 2016](#)). In Ethiopia, the prevalence of low birth weight varies from place to place. It was reported 28.3% in Kersa district, Oromia region, eastern Ethiopia ([Assefa, Berhane, & Worku, 2012](#)) and 22.5% in Jimma zone, Southwest Ethiopia ([Conde-Agudelo & Díaz-Rossello, 2016](#)), 17.4% in Gonder, northern Ethiopia ([Tadesse, 2014](#)).

The implementation of KMC also focuses on the decision-making process not depends only on the mother's desire and willingness, but also on the support provided by the family members as well as kindhearted health care teams ([Chisenga, Chalanda, & Ngwale, 2015](#)).

Study showed that only about 10% to 25% of

preterm and/or low birth weight babies receive KMC in Ethiopia between 2015–2017, the estimates was derived from an assessment that was conducted in two hospitals that were providing KMC services in Ethiopia, after training; these hospitals provided KMC services to 36% of preterm babies ([Lakew & Worku, 2014](#)). Mothers are the main caregiver to newborns hence the care is habitually dependent on their level of knowledge, attitude, and practice (KAP) about newborn care. Therefore, this study aimed to study the KAP of mothers about KMC.

METHODS

Study Setting and Participants

Cross-sectional study was conducted from 1 March to 30 March 2020 in HiwotFana Specialized University hospital located in Harar, Ethiopia. Harar is located at eastern direction of 526 km away from the capital Addis Ababa. HiwotFana Specialized University hospital is one of the specialized university hospitals in the country which was established in 2006 and providing service for more than 5 million people in the catchment area.

Sample Size Determinations and Sampling Techniques

Sample size was calculated by using single population proportion formula by taking 19.53 % from a similar study ([Jamie & Ahmed, 2020](#)). Using the 95% CI and 5% marginal error (d) ($n = (Z_{\alpha/2})^2 p(1-p) / d^2$), the given sample size is 242. According to Hiwotfana specialized university Hospital's delivery report, a total of 400 women delivered per month. Since the total population is less than 10,000, correction formula was applied ($n = n / (1 + n/N)$). So, the final sample size after adding 10% non-respondent rates was 166.

HiwotFana Specialized university hospital provides advanced services including Neonatal Intensive Care Unit (NICU) and KMC units, so the hospital was chosen deliberately due to this reason. Preterm and/or low birth weight (LBW) who were born among postnatal mothers were selected successively until the needed sample size was obtained. If there were two or more one baby was selected by lottery method to the study.

Normal postnatal mothers having pre term, LBW, stable infants breathing on their own, and infants without life-threatening disease or malformations that are willing to participate were included in this study.

Ethical Clearance

Ethical clearance was obtained from Ethical Clearance Committee of Harar Health Science College (Reference number: HHSC-093/2020). Consent was obtained from the administrative bodies of the health facility and from the participants.

Data Collection and Data Analysis

Data were collected by face-to-face interview using standardized questionnaire. For practice part, direct observation was done when they position, breast feed and ambulate. After data collection, the questionnaire was checked for completeness and coded. The data were entered into Epi-info version 3.5.3 and exported, cleaned and analyzed by using SPSS version-20. Descriptive analyses were performed and

bivariate analyses were used to find out the association of independent variables. Variables with a $p < 0.05$ in the bivariate analysis were entered into multiple logistic regression and variables with $p < 0.05$ in the multivariate analysis were considered to have statistically significant associations.

RESULTS

Socio-Demographic Characteristics

One hundred sixty-six (166) postnatal mothers with preterm /low birth weight were participated in this study 97 (58.43%) were age 18-35 years, 96 (57.83%) were residing in rural area, more than half 90 (54.22%) were illiterate, 146 (87.95%) of the neonates included in this study were healthy. Near to all 158 (95.18%) had ANC follow up. Majority 97 (58.43%) of deliveries was spontaneous vaginal deliveries (SVD). Majority of them 89 (53.61%) were preterm by gestational age and nearly all 162 (97.59%) were LBW by birth weight group (See Table 1).

Table 1 Socio demographic characteristics of participants in a tertiary care center of Harari region, Harar, Ethiopia, 2020

Variable		Frequency	(%)
Age group (years)	18-35	97	58.43
	> 35	69	41.57
Marital status	Single	5	3.01
	Married	107	64.46
	Widows	7	4.22
	Divorced	47	28.31
Residence	Urban	70	42.17
	Rural	96	57.83
ANC follow up	Yes	158	95.18
	No	8	4.82
Level of formal education	No formal education	90	54.22
	Primary	49	29.52
	Secondary	19	11.45
	Tertiary	8	4.82
Sex of infant	Male	78	46.99
	Female	88	53.01
Health status	Healthy	146	87.95
	Sick	20	12.05
Type of delivery	SVD	97	58.43
	Instrumental	14	8.43
	C/S	55	33.13
Type of birth	Singleton	157	94.58
	Multiple	9	5.42
Gestational age	Preterm	89	53.61
	Term	77	46.39
Birth weight	LBW	162	97.59
	Very LBW	3	1.81
	Extremely LBW	1	0.60

Knowledge of Care Takers about the KMC

Overall study participants 137 (82.53%) had heard about kangaroo mother care (KMC). Among those, all 137 (82.53%) of respondents answered it as skin to skin contact of the infant on the mother's chest, whereas the rest of them did not know. One hundred thirty-seven (82.53%) caregivers know at least one benefit of KMC.

Study subjects were asked to their knowledge on the importance of KMC: 137 (82.53%) provides warmth to the baby, 117 (70.48%) promotes

exclusive breast feeding, 93 (56.02%) improves weight gain and growth, 71 (42.77%) reduces hospital stay, 93 (56.02%) reduces infection and 137 (82.53%) promotes baby-mother bonding. Regarding appropriate initiation time to KMC 93 (67.88%) of the answered that it should be started immediately after birth (See **Table 2**). The majority of the participants respond that the heard the information about kangaroo mother care through health professionals, mass media, friends, and posters 86%, 65%, 23.3% and 17.4%, respectively.

Table 2 Knowledge towards KMC Among Postnatal Mothers in A Tertiary Care Centre of Harari Region, Harar, Ethiopia, 2020

Knowledge towards KMC		n (%)
Do you hear about KMC	Yes	137 (82.53%)
	No	29 (17.47%)
What is KMC	Skin to skin contact to stabilizes baby's temperature	137 (82.53%)
	Incubator care stabilizes baby's temperature	29 (17.47%)
KMC provides warmth to the baby	Yes	137 (82.53%)
	No	29 (17.47%)
KMC promotes bonding between mother and baby	Yes	137 (82.53%)
	No	29 (17.47%)
KMC improves the mother's confidence in handling her baby	Yes	87 (52.41%)
	No	79 (47.59%)
KMC promotes breast feeding	Yes	117 (70.48%)
	No	49 (29.52%)
KMC promotes mental development of premature babies.	Yes	82 (49.40%)
	No	84 (50.60%)
Babies who are given kangaroo care sleep deeply and wake up less often	Yes	97 (58.43%)
	No	69 (41.57%)
KMC prevents postpartum depression	Yes	99 (59.64%)
	No	67 (40.36%)
Babies are who given KMC cry less	Yes	124 (74.70%)
	No	42 (25.30%)
KMC promotes baby's growth and development	Yes	93 (56.02%)
	No	73 (43.98%)
KMC results in reduced infection in the baby	Yes	93 (56.02%)
	No	73 (43.98%)
KMC leads to early discharge	Yes	71 (42.77%)
	No	95 (57.23%)

Attitudes of Care Takers on KMC

One hundred thirty-seven (82.53%) of mothers believed positive regarding implementation of KMC for it corrects the temperature and promotes growth and development of their small babies. On the contrary, only 19 (11.45%)

respondents answered that both parents should be involved in kangaroo care, the majority left 147 (88.55%) said that only mother should be involved for the care of the neonates (See **Table 3**).

Table 3 Attitude towards KMC Among Postnatal Mothers in A Tertiary Care Centre of Harari Region, Harar, Ethiopia, 2020

Variable	Agree	Not agree
KMC provides warmth to the baby	137 (82.53%)	29 (17.47%)
KMC promotes bonding between mother and baby	137 (82.53%)	29 (17.47%)
KMC improves the mother's confidence in handling her baby	87 (52.41%)	79 (47.59%)

Table 3 (Cont.)

KMC promotes breast feeding	117 (70.48%)	49 (29.52%)
KMC promotes mental development of premature babies.	82 (49.40%)	84 (50.60%)
Both parents should be involved in kangaroo care.	19 (11.45%)	147 (88.55%)
Babies who are given kangaroo care sleep deeply and wake up less often	97 (58.43%)	69 (41.57%)
Incubator care stabilizes baby's temperature than KMC	29 (17.47%)	137 (82.53%)
KMC prevents postpartum depression	99 (59.64%)	67 (40.36%)
Babies are who given KMC cry less	124 (74.70%)	42 (25.30%)
KMC promotes baby's growth and development	137 (82.53%)	29 (17.47%)
KMC results in reduced infection in the baby	93 (56.02%)	73 (43.98%)
KMC leads to early discharge	71 (42.77%)	95 (57.23%)
KMC should be started immediately after birth	94 (56.63%)	72 (43.37%)

Practice of Care Takers on KMC

Participants were asked and observed on the correct practice of Kangaroo mother care on kangaroo position, clothing during KMC, KMC initiation and type of KMC. About 44 (32.12%) of respondents performed kangaroo position in setting or semi-reclined position appropriately, whereas the rest of respondents' position was inappropriate. Most 49 (57%) of respondents

dressed the baby with cap, socks and nappy appropriately, whereas the rest of them did not dress their baby properly. Nearly all 123 (89.70%) of KMC was performed by mothers, 14 (10.22%) were performed by others which include grandmothers, aunts and maids, but there was no single one which was performed by fathers (See **Table 4**).

Table 4 Practice Towards KMC Among Postnatal Mothers in A Tertiary Care Centre of Harari Region, Harar, Ethiopia, 2020

Practice Towards KMC		n	%
KMC was performed by	Mother	123	89.7
	Father	0	0.00
	Others	14	10.22
Position for KMC	Proper	44	32.12
	Not proper	93	67.88
Dressing during KMC	Dressed cap, sock& nappy	44	32.12
	No dressed well	93	67.88
Type of KMC	Continuous	86	62.77
	Intermittent	51	37.23

Over all knowledge, attitude and practice score sum of the care givers towards KMC illustrated as follow.




Figure 1 Knowledge, Attitude and Practice towards KMC Among Postnatal Mothers in A Tertiary Care Centre of Harari Region, Harar, Ethiopia

DISCUSSION

Findings of this study revealed the level of good knowledge among mothers (82.53%), which was very high compared to the study done in Rohilkhand Hospital Bareilly that the knowledge was 33.3% ([Sharma & Verma, 2019](#)). Whereas it was lower than the study done in North Ethiopia (99.3%) ([Tsegay, 2015](#)), Malawi (84%) ([Chisenga et al., 2015](#)), and Ghana (90%) ([Nguah et al., 2011](#)). This difference might be due to a difference in socio-demographic/cultural /economic characteristics in different regions and methodological difference.

This study showed that 82.53% of mothers had positive attitude towards KMC. The finding was lower than the study done among Indian mothers, which the acceptance of KMC was 96% ([Parmar et al., 2009](#)), and the study done in Cape Town with the acceptance of KMC was 96.6%, and 87.1% mothers who had favorable opinion towards the practice of KMC ([Rosant, 2009](#)). This could be influenced by socio-demographic/cultural /economic characteristics in different regions and methodological difference.

In this study, the level of correct practice was 32.12%, it is lower than a study done at Hawasa University Hospital which was 61.6% ([Getinet K., 2008](#)). This difference may be due to the cultural characteristics of the respondents, study methodology and set up of services. Also, 42.77% of the participants agreed that KMC reduces hospital stay, the result was lower than that of a study done in South India 56.69% strongly agreed that KMC reduces hospital stay ([Suhasimi Mekala, 2019](#)). This difference could be related to the difference in study methodology.

Our findings also indicated that the participants reported the benefits of KMC, such as protection from infection, warmth to the baby, exclusive breast feeding, weight gain and growth, early discharge from hospital, safety and love. These have been also reported by other studies ([March of Dimes, PMNCH, Save the Children, & WHO, 2012](#); [WHO, 2003](#)).

Since the study incorporates only public facility but not private facilities, so the findings may be difficult generalizable for general population.

CONCLUSION

The finding of this study on knowledge, attitude and practice of caretakers regarding kangaroo mother care showed that 82.53 % had good knowledge, 82.53 % had positive attitude, and 32.2 % practiced correctly. However, there were clear gaps in knowledge and practice in the study. Therefore, it is very important to have health education sessions during ANC follow-up on KMC. Furthermore, it is recommended to conduct qualitative research to explore the temporal relationship.

Declaration of Conflicting Interest

The author declares no conflict of interest.

Funding

The author declares that no financial support for this research.

Acknowledgements

The author would like to express his gratitude to the Harar Health Science College and Hiwot-Fana Specialized Referral Hospital, NICU, KMC unit and registration room heads of the hospital and data collectors.

References

- Akter, T., Dawson, A., & Sibbritt, D. (2016). What impact do essential newborn care practices have on neonatal mortality in low and lower-middle income countries? Evidence from Bangladesh. *Journal of Perinatology*, 36(3), 225-230.
- Assefa, N., Berhane, Y., & Worku, A. (2012). Wealth status, mid upper arm circumference (MUAC) and antenatal care (ANC) are determinants for low birth weight in Kersa, Ethiopia. *PloS one*, 7(6), e39957.
- Chisenga, J. Z., Chalanda, M., & Ngwale, M. (2015). Kangaroo Mother Care: A review of mothers' experiences at Bwaila hospital and Zomba Central hospital (Malawi). *Midwifery*, 31(2), 305-315.
- Conde-Agudelo, A., & Diaz-Rossello, J. L. (2016). Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. *Cochrane Database of Systematic Reviews*(8).
- Getinet K. (2008). Knowledge, attitude, practice and associated factors of kangaroo mother care for neonatal survival among care takers of preterm and low birth weight infants in health care settings, Hawassa, Southern Ethiopia, *Journal of Health, Medicine and Nursing*, 58: 1-9.
- Jamie, A. H., & Ahmed, A. M. (2020). Factors associated with low birth weight among newborns in Ethiopia. *Public Health of Indonesia*, 6(1), 1-6.

- Lakew, W., & Worku, B. (2014). Follow-up profile and outcome of preterms managed with kangaroo mother care. *Open Journal of Pediatrics*, 4(2), 143-147.
- Lawn, J. E., Cousens, S., Zupan, J., & Lancet Neonatal Survival Steering Team. (2005). 4 million neonatal deaths: when? Where? Why?. *Lancet (London, England)*, 365(9462), 891-900.
- M Ludington-Hoe, S. (2011). Evidence-based review of physiologic effects of kangaroo care. *Current Women's Health Reviews*, 7(3), 243-253.
- March of Dimes, PMNCH, Save the Children, & WHO. (2012). *Born too soon: The global action report on preterm birth* (CP Howson, MV Kinney, & J. Lawn Eds.). Geneva: World Health Organization.
- Nguah, S. B., Wobil, P. N., Obeng, R., Yakubu, A., Kerber, K. J., Lawn, J. E., & Plange-Rhule, G. (2011). Perception and practice of Kangaroo Mother Care after discharge from hospital in Kumasi, Ghana: A longitudinal study. *BMC Pregnancy and Childbirth*, 11(1), 99.
- Nyqvist, K. H., Anderson, G., Bergman, N., Cattaneo, A., Charpak, N., Davanzo, R., . . . Mendoza, S. (2010). Towards universal kangaroo mother care: Recommendations and report from the first European conference and seventh international workshop on kangaroo mother care. *Acta Paediatrica*, 99(6), 820-826.
- Parmar, V. R., Kumar, A., Kaur, R., Parmar, S., Kaur, D., Basu, S., . . . Narula, S. (2009). Experience with Kangaroo mother care in a neonatal intensive care unit (NICU) in Chandigarh, India. *The Indian Journal of Pediatrics*, 76(1), 25-28.
- Rajaratnam, J. K., Marcus, J. R., Flaxman, A. D., Wang, H., Levin-Rector, A., Dwyer, L., . . . Murray, C. J. (2010). Neonatal, postneonatal, childhood, and under-5 mortality for 187 countries, 1970–2010: a systematic analysis of progress towards Millennium Development Goal 4. *The Lancet*, 375(9730), 1988-2008.
- Rosant, C. (2009). *Knowledge of and attitudes towards kangaroo mother care in the Eastern Subdistrict, Cape Town*. Cape Town: University of the Western Cape.
- Sharma, K., & Verma, V. L. (2019). A descriptive study to assess the knowledge regarding kangaroo mother care among postnatal mother with the view to develop information pamphlets in Rohilkhand Medical College and Hospital Bareilly, *International Journal of Advance Research, Ideas and innovations in Technology*, 5(3), 1462-1466.
- Suhasimi Mekala. (2019). Kangaroo mother care: Knowledge, attitudes and practices among nursing mothers and its impact on weight gain in low birth weight babies. *Indian Journal of Applied Research*, 9(9), 57-60.
- Taddesse, K. (2014). A low birth weight and associated factors among newborns in Gonder town. *Indo Global Journal of Pharmaceutical Science*, 4, 74-80.
- Trevisanuto, D., Putoto, G., Pizzol, D., Serena, T., Manenti, F., Varano, S., . . . Wingi, O. (2016). Is a woolen cap effective in maintaining normothermia in low-birth-weight infants during kangaroo mother care? Study protocol for a randomized controlled trial. *Trials*, 17(1), 1-6.
- Tsegay, T. (2015). *Knowledge and practice of immediate newborn care among midwives in governmental health facilities at Central Zone, Tigray Regional State, North Ethiopia*. Ethiopia: Addis Ababa University.
- WHO. (2003). *Kangaroo mother care: A practical guide*. France: World Health Organization.
- WHO. (2017). *Neonatal mortality*. Retrieved from http://www.who.int/gho/child_health/mortality/neonatal/en/index1.html
- WHO, & UNICEF. (2004). *Low birthweight: Country, regional and global estimates*. Retrieved from <https://apps.who.int/iris/handle/10665/43184>
- WHO, & UNICEF. (2009). *WHO-UNICEF Joint Statement on home visits for the newborn child: A strategy to improve survival*. Retrieved from Geneva: https://www.who.int/maternal_child_adolescent/documents/who_fch_cah_09_02/en/

Cite this article as: Jamie, A. H. (2020). Knowledge, attitude and practice of kangaroo mother care among postnatal mothers in a tertiary care center of Harari region, Harar, Ethiopia. *Public Health of Indonesia*, 6(4), 116-122. <https://dx.doi.org/10.36685/phi.v6i4.355>