

Prevalence and associated factors of Smokeless Tobacco (SLT) use among adolescents in Indonesia: GYTS 2019 data analysis

Syamsuriansyah^{1*}, Salman², Heni Rusmitasari³, Marlynda Happy Nurmalita Sari⁴, and Maretalinia⁵

¹Department of Medical Record, Politeknik Medica Farma Husada Mataram, Indonesia

²Department of Pharmacy, Faculty of Health Science, Singaperbangsa Karawang University, Indonesia

³Fakultas Kesehatan Masyarakat, Universitas Muhammadiyah Semarang, Indonesia

⁴Department of Midwifery, Poltekkes Kemenkes Semarang, Indonesia

⁵Institute for Population and Social Research, Mahidol University, Thailand

Doi: <https://dx.doi.org/10.36685/phi.v8i1.569>

Received: 25 February 2022 | Revised: 5 March 2022 | Accepted: 7 March 2022

Corresponding author:

Syamsuriansyah

Departement of Medical Record, Politeknik Medica Farma Husada Mataram

Jl. Medica Farma No.1, Baturungit Selatan, Kec. Sekarbela, Kota Mataram, Indonesia 83116

Email: sam_bptk@yahoo.com

Copyright: © 2022 the Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium provided the original work is properly cited.

Abstract

Background: Smokeless tobacco (SLT) use is a new trend since the banning of smoking in many public places, including among adolescents. Many people believe that SLT is a harmful reduction method compared to cigarettes. However, there are many risks of using the SLT, such as Head and neck cancer (HNC), cancer of the mouth, and coronary heart disease.

Objective: This study aimed to examine the prevalence and the factors associated with SLT use among adolescents in Indonesia.

Methods: This study used the cross-sectional data from the Global Youth Tobacco Survey (GYTS) Indonesia 2019. The national school was selected by proportional random sampling. Around 8,828 students were eligible for this study.

Results: This study found the prevalence of SLT use was 3.25%. The bivariate test found each independent variable was significantly associated with the SLT use. However, the adjusted test found that age, smoking parent, smoking teacher, knowledge, and attitude were significantly associated with SLT use.

Conclusion: The variables mostly related to SLT use come from the family and school environment. The appropriate intervention needs to be arranged to reduce the prevalence of smokers, including SLT use.

Keywords: Smokeless tobacco (SLT); adolescent; Global Youth Tobacco Survey (GYTS); Indonesia

Background

Smokeless tobacco (SLT) is an alternative way of smoking, and the reduction of the effect is still debated ([Centers for diseases control and](#)

[prevention, 2020](#)). SLT is the product in which the nicotine is put in the mouth so the nicotine will be absorbed through the lining of the mouth ([Ebbert et al., 2015](#)). Nicotine is the most ingredient containing tobacco, including SLT. The result of one

study found that nicotine addiction and millions of deaths per year worldwide increase due to cancer (Hecht & Hatsukami, 2022). In addition, the Centers for Disease Control and Prevention (CDC) reported the health problems that have an association with smokeless tobacco control, such as nicotine addiction, cancer of the mouth, esophagus, and pancreas, diseases of the mouth, early delivery, and stillbirth during pregnancy, nicotine poisoning in the children, and increase the risk for death from heart diseases and stroke (Centers for diseases control and prevention, 2020). There are several smokeless products such as snus, chew, dry snuff, moist snuff, and dissolvable (Delnevo et al., 2014; Skulberg et al., 2019).

The trend of smoke in developing countries, including Indonesia, is increasing since the cigarette price in Indonesia is very cheap, which was around 20.830 IDR on average in 2020 (Statista, 2020). A low tax has been applied, which leads to a higher number of new smokers (Prasetyo & Adrison, 2020). Compared with other countries, the tax for cigarettes in Indonesia is very low (Prasetyo & Adrison, 2020). The affordability of the student to purchase cigarettes is very dependent on the number of money pocketed (Saari et al., 2014). The higher number of Non-Communicable Diseases (NCD) since the phenomena is starting high from the epidemiological transition (Sylvestre et al., 2018). Being a passive smoker increases the probability of having NCD, which is higher than smoker (World Health Organization, 2019). As an alternative to reducing the health risk, some communities decided to use smokeless tobacco. However, the SLT does not exactly reduce the harmful ingredients because the nicotine is put in the mouth, so the risk of cancer of the mouth is high (Jalilian et al., 2015; Suryati & Tarigan, 2019; Wibowo et al., 2019). Another study using Indonesia GYTS (Global Youth Tobacco Survey) 2014 revealed that friends, parents, and teachers are the role model to influence a student to be a smoker (Jamal et al., 2020)

GYTS is a global project managed by World Health Organization. The Indonesia GYTS is designed to estimate the worldwide burden of tobacco use among adolescents (Eriksen et al., 2012). In Indonesia, the number of tobacco users in adolescent school age always increases. It

increased a lot from 0.3% in 1995 to 2.0% in 2007 (Jalilian et al., 2015; Suryati & Tarigan, 2019). The data from Basic Health Research Indonesia in 2013, the prevalence of smokers among adolescents was 18.3% (Kementerian Kesehatan Republik Indonesia, 2018). Comparing the place of residence, in 2017, smokers in rural (1.55%) are higher than in urban (1.07%) (Syachroni et al., 2013). The main findings from GYTS 2019 revealed that 19.2% of students are using tobacco, but around 1% of them currently use smokeless tobacco products (Windiarto et al., 2018). This study focuses on smokeless tobacco use to find the prevalence and associated factors of SLT among adolescents using GYTS 19 data.

Methods

Study Design

The original survey, GYS 2019, is a cross-sectional study conducted in 34 provinces in 2019. GYTS aimed to control tobacco use among adolescents aged 13 to 17 years (World Health Organization, 2020). The current study aimed to explore the prevalence of SLT use and factors associated with SLT use in adolescents using GYTS 2019 data.

Samples/Participants

The original survey selected the school based on proportional probability in terms of the sampling method. The class was chosen by random sampling method, and all the eligible students in grades 7-12 could join the survey. This study used the sample who were eligible for this study and met the inclusion and exclusion criteria. The inclusion criteria are a student aged 7 to 12 years old. The exclusion criteria are those who complete the questionnaire. The total sample in the original survey was 8,823 students.

Instruments

This current study used the smokeless tobacco (SLT) use-dependent variable. It was a dummy variable with categories *yes/no*. The independent variables used in this study were the sociodemographic variables, knowledge, and attitude. The question about tobacco use (SLT use) was chosen due to the prevalence of cigarette use decreasing in some communities, but on another side, the SLT use is increasing. The Sociodemographic questions were taken from the original questionnaire. Those independent variables

were chosen to examine the correlation of the predictors to the SLT use. The study about determinants of SLT use is important since the lack of studies discuss SLT use compared to the common cigarette use.

Data Analysis

The statistical analysis used consists of univariate, bivariate, and multivariate. The univariate showed the frequency and percentage (for categorical variable) and minimum number, maximum number, and mean (for continuous variable) of each variable to see the respondents' characteristics. The bivariate analysis with Chi-Square and t-test to see the Crude Odd Ratio (COR) for each independent variable to dependent variables. The binary logistic regression was done to examine all the independent variables to the dependent variables in three models, which were presented by Adjusted Odd Ratio (AOR). STATA version 15 Mahidol University licensed was used to analyze the data.

Results

About 8,823 respondents were eligible for this study. Among them, about 3.25% used smokeless tobacco (SLT), which means cigarettes with low nicotine. In terms of the age of students, most of them were in age 13 to 14 years old. More than half of the respondents were female, and most of the students were in grades 1 to 3. According to the money pocket, more than half of the respondents had more than 20,000 rupiahs per day. Many of their parents do not smoke; as shown by the data, 53% of the parents were not smokers. In the school environment, most of the teachers were smokers, as demonstrated by the data that many teachers smoke during school hours (**Table 1**). In terms of knowledge and attitude, most of the respondents have known that smoke from others is dangerous to their health, and they agree with the policy of banning smoking in any enclosed public place.

The bivariate analysis using the Chi-square test found all independent variables significantly associated with the smokeless tobacco use, except the grade, which showed a low correlation. After adjusted analysis, the binary logistic regression test result found that higher age tends to decrease the tendency to be smokeless tobacco users compared to the young one. Compared with the students having no smoker parents, students having mother

smokers significantly increase the likelihood 2.22 times for using the smokeless tobacco. In the school environment, compared with teachers who smoke every day during school hours, teachers who never smoke during the school hours decreased the probability of students using smokeless tobacco. In terms of the knowledge which showed by the questions of harmful smoke from others, compared to the students who thought the smoke from the other cigarettes is not harmful, those who thought it was harmful decrease the probability of smoking the smokeless tobacco. Compared to students who agree with banning smoking in any enclosed public place, those who did not agree were 1.4 times more likely to use smokeless tobacco. However, the variables of sex, grade, and money pocket did not correlate with smokeless tobacco use after testing the binary logistic regression (**Table 1**).

Discussion

The prevalence of smokeless tobacco (SLT) use in Indonesia remains low. However, the prevalence is going to elevate since there are many policies banning smoke in many places. Another study using IFLS (Indonesia Family Life Survey) 2015 wave five found the prevalence of SLT use was 5.6% ([Pengpid & Peltzer, 2019](#)). The SLT may contain harmful ingredients. One study reported that SLT might contain metals such as arsenic ([Muthukrishnan & Warnakulasuriya, 2018](#)). Another study in India revealed that consuming the SLT may have the risk of cancer, especially oral cancer ([Boffetta et al., 2005](#)). In line with the previous study, one study found SLT use correlated to Head and Neck Cancer (HNC), especially oral cancers ([Wyss et al., 2016](#)). Different results were shown from another study that mentioned that SLT use is not correlated to cancer but is mostly correlated to coronary heart disease ([Timberlake et al., 2017](#)). SLT products are increasing as the campaign of non-smoking and banning smoke in the enclosed public place ([Delnevo et al., 2014](#)). For instance, the moist snuff products increased 65.6% from 2005 to 2011 ([Delnevo et al., 2014](#)). SLT products are not the harm reduction method compared to cigarettes, and further research is needed to explore that ([Hatsukami et al., 2004](#)). The study examined the determinants of SLT in 140 countries found that the higher prevalence of SLT was in low- and lower-income countries ([Sinha et](#)

al., 2011). The number of SLT users in Sri Lanka is increasing since the low economic status community tends to have low awareness about the health risk of using SLT (Somatunga et al., 2012). The recent study revealed the factors influencing the SLT use were gender, educational level, wealth index, place

of residence, socioeconomic status, and low tax policy (Thakur & Paika, 2018). Another study found that factors such as former smoking status, low education level, living in rural areas, younger age, white race, and unemployment were associated with SLT use (Chang et al., 2016).

Table 1 Characteristic of the respondents, Chi-square, and binary logistic regression results

Variables (N = 8,823)	n (%)	Smokeless tobacco use	
		Chi-Square	AOR. CI 95%
Tobacco Use			
Used smokeless tobacco (Yes)	287 (3.25)		
Sociodemographic Factors			
How old are you?		0.000	
11 years old or younger	247 (2.80)		Ref
12 years old	1,156 (13.10)		0.50* (0.28-0.90)
13 years old	1,530 (17.34)		0.33*** (0.18-0.60)
14 years old	1,575 (17.85)		0.50* (0.29-0.88)
15 years old	1,435 (16.26)		0.44** (0.24-0.80)
16 years old	1,418 (16.07)		0.62 (0.32-1.19)
17 years old	1,462 (16.57)		0.68 (0.36-1.28)
Sex		0.000	
Male	3,677 (41.68)		Ref
Female	5,146 (58.32)		1.31 (0.95-1.81)
Grade		0.010	
Grade 1-3	5,074 (57.51)		Ref
Grade 4-6	3,749 (42.49)		0.69 (0.46-1.03)
During an average week, how much money do you have that you can spend on yourself; however you want?		0.000	
Less than Rp. 20,000	4,025 (45.62)		Ref
More than Rp. 20,000	4,798 (54.38)		0.79 (0.61-1.02)
Do your parents smoke tobacco?		0.000	
None	4,651 (52.71)		Ref
Father only	3,422 (38.78)		0.82 (0.62-1.08)
Mother only	84 (0.95)		2.22*(1.11-4.48)
Both	249 (2.82)		1.29 (0.74-2.27)
Don't know	417 (4.73)		1.03 (0.60-1.77)
Teacher smoking during school hours		0.000	
About every day	941 (10.67)		Ref
Sometimes	3,150 (35.70)		1.80 (0.755-1.54)
Never	3,120 (35.36)		0.64* (0.42-0.95)
Do not know	1,612 (18.27)		0.068 (0.43-1.07)
Knowledge and Attitudes			
Do you think the smoke from other people's cigarettes is harmful to you?		0.000	
Definitely not	250 (2.83)		Ref
Probably not	147 (1.67)		1.38 (0.69-2.75)
Probably yes	1,665 (18.87)		0.45** (0.26-0.78)
Definitely yes	6,761 (76.63)		0.49** (0.29-0.82)
Are you in favor of banning smoking inside enclosed public places?		0.000	
Yes	7,936 (89.95)		Ref
No	887 (10.05)		1.40* (1.00-1.96)

*p-value < 0.05, **p-value < 0.01, ***p-value < 0.001

Number of obs = 8,823 | LR chi2(40) = 251.82 | Prob >chi2 = 0.0000 | Pseudo R2 = 0.0995 | Log likelihood = -1139.53

The limited study examined the smokeless tobacco (SLT) use in Indonesia leads to elaborate the explanation through the general cigarettes. The prevalence of current tobacco use among adolescents in Indonesia remains high compared with the previous survey, 19.2% (35.6% male and 3.5% female) ([World Health Organization, 2020](#)). The prevalence of tobacco use in Indonesia was higher than in Cameroon, Myanmar, Botswana, and India, with current adolescent smoking was 11.2%, 13.6%, 10%, and 11.4%, respectively ([Lalithambigai et al., 2016](#); [Mbatchou Ngahane et al., 2015](#); [Mbongwe et al., 2017](#); [Tun et al., 2017](#)). Compared to 10 countries in Africa, the current cigarette smoking among students aged 13-15 was in the range of 3.4% to 13.6%, lower than in Indonesia ([Zhao et al., 2016](#)). The access and affordability to purchase cigarettes near the school may be due to the kiosk or shop near the school selling tobacco products. The knowledge of cigarettes might come from the family who spends the majority of their time with the adolescent ([Maretalinia et al., 2021](#)). This study found that the SLT is determined by age, parents smoking, teacher smoke, knowledge, and attitude.

The regulation from the Minister of Education mentioned that the school environment is a free smoke area that needs to be supported by the headmaster of the school, teachers, students, and other people in the school area ([Ministry of Education Republic of Indonesia, 2015](#)). School is prohibited from producing, selling, distributing, sponsoring, and putting a cigarettes logo. The regulation about addictive products also mentioned that smoking cigarettes are prohibited in public areas, including schools. The regulation of the Minister of Finance mentioned the price of cigarette sticks ([Ministry of Finance Republic of Indonesia, 2021](#)). The cigarette advertisement is prohibited from demonstrating by using the real cigarette, and there is a time restriction of promotion in television ([Barker et al., 2019](#); [Bigwanto & Soerojo, 2020](#); [Blencowe et al., 2016](#); [Lyons et al., 2014](#); [World Health Organization, 2013](#)).

Smokeless tobacco (SLT) use is one solution to reduce the negative effect of tobacco on health. However, that statement is still debated. The determinants of SLT use can give the general characteristics of SLT users among adolescents

and the factors that influence them to choose SLT compared to other types of tobacco. In the public health area, the study about SLT is not many, especially the study using the national representative data among adolescents. Therefore, the government, especially the Ministry of Education, needs to collaborate with the Ministry of Health to arrange the program to reduce the prevalence of tobacco use, especially SLT use. Furthermore, the appropriate planning programs may prevent adolescents from smoking. For instance, the communication, information, and education program using the interesting application with adolescent-friendly. That application in the smartphone can provide information about the harmfulness of smoking, the diseases caused by smoking, and tips on how to get great achievement as a student without smoking.

Conclusion

The prevalence of smokeless tobacco (SLT) use among adolescents in Indonesia in 2019 was 3.25%. The associated factors of SLT use were mostly related, such as age, smoking parents, smoking teacher, knowledge, and attitude. Multilevel prevention and control can reduce the prevalence of SLT use among adolescents, starting from a student level, school level, household level, community level, and national level. There is a need the communication, information, and education about the harmfulness of the SLT to change the perspective of risk reduction by using SLT.

Declaration of Conflicting Interest

All of the authors declare no conflict of interest in this study.

Funding

None.

Acknowledgment

Thank you for Center for Disease Control and Prevention (CDC) was provided open this data online.

Author Contribution

All authors contributed to all phases of the study and concurred with the last version of the article to be issued and accountable in all stages of the work.

Author Biographies

DR. Syamsuriansyah is a Director of Politeknik Medica Farma Husada, Mataram, Indonesia.

Salman is a Junior Lecturer at the Singaperbangsa Karawang University, Indonesia.

Heni Rusmitasari is a Lecturer at Faculty of Public Health, Universitas Muhammadiyah Semarang, Indonesia.

Marlynda Happy Nurmalita Sari is a Lecturer at Poltekkes Kemenkes Semarang, Indonesia.

Maretalinia is a Ph.D. Student at Institute Population and Social Research, Mahidol University, Thailand.

References

- Barker, A. B., Whittamore, K., Britton, J., & Cranwell, J. (2019). Content analysis of tobacco content in UK television. *Tobacco Control*, 28(4), 381-385. <https://doi.org/10.1136/tobaccocontrol-2018-054427>
- Bigwanto, M., & Soerojo, W. (2020). Content analysis of cigarette packs in Indonesia: Regulatory non-compliance and product promotion advantage. *Public Health Indonesia*, 6, 18-27. <https://doi.org/10.36685/phi.v6i1.330>
- Blencowe, H., Cousens, S., Jassir, F. B., Say, L., Chou, D., Mathers, C., Hogan, D., Shiekh, S., Qureshi, Z. U., You, D., & Lawn, J. E. (2016). National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *The Lancet Global Health*, 4(2), e98-e108. [https://doi.org/10.1016/s2214-109x\(15\)00275-2](https://doi.org/10.1016/s2214-109x(15)00275-2)
- Boffetta, P., Aagnes, B., Weiderpass, E., & Andersen, A. (2005). Smokeless tobacco use and risk of cancer of the pancreas and other organs. *International Journal of Cancer*, 114(6), 992-995. <https://doi.org/10.1002/ijc.20811>
- Centers for Diseases Control and Prevention. (2020). Smokeless tobacco: health effects. USA: Centers for Diseases Control and Prevention
- Chang, J. T., Levy, D. T., & Meza, R. (2016). Trends and factors related to smokeless tobacco use in the United States. *Nicotine and Tobacco Research*, 18(8), 1740-1748. <https://doi.org/10.1093/ntr/ntw090>
- Delnevo, C. D., Wackowski, O. A., Giovenco, D. P., Manderski, M. T., Hrywna, M., & Ling, P. M. (2014). Examining market trends in the United States smokeless tobacco use: 2005-2011. *Tobacco Control*, 23(2), 107-112. <https://doi.org/10.1136/tobaccocontrol-2012-050739>
- Ebbert, J. O., Elrashidi, M. Y., & Stead, L. F. (2015). Interventions for smokeless tobacco use cessation. *Cochrane Database of Systematic Reviews*, 2015(10), Cd004306. <https://doi.org/10.1002/14651858.CD004306.pub5>
- Eriksen, M., Mackay, J., & Ross, H. (2012). *The tobacco atlas* (4th ed.). Atlanta, GA, USA: American Cancer Society.
- Hatsukami, D. K., Lemmonds, C., & Tomar, S. L. (2004). Smokeless tobacco use: Harm reduction or induction approach? *Preventive Medicine*, 38(3), 309-317. <https://doi.org/10.1016/j.ypmed.2003.10.006>
- Hecht, S. S., & Hatsukami, D. K. (2022). Smokeless tobacco and cigarette smoking: chemical mechanisms and cancer prevention. *Nature Reviews Cancer*, 22(3), 143-155. <https://doi.org/10.1038/s41568-021-00423-4>
- Jalilian, F., Karami Matin, B., Ahmadpanah, M., Ataee, M., Ahmadi Jouybari, T., Eslami, A. A., & Mirzaei Alavijeh, M. (2015). Socio-demographic characteristics associated with cigarettes smoking, drug abuse and alcohol drinking among male medical university students in Iran. *Journal of Research and Health Sciences*, 15(1), 42-46.
- Jamal, H., Abdullah, A. Z., & Abdullah, M. T. (2020). Determinan sosial perilaku merokok pelajar di Indonesia: analisis data global youth tobacco survey tahun 2014. *Jurnal Kesehatan Vokasional*, 5(3), 141-150. <https://doi.org/10.22146/jkesvo.56718>
- Kementerian Kesehatan Republik Indonesia. (2018). *Hasil utama RISKESDAS 2018*. Jakarta: Kemenkes RI.
- Lalithambigai, G., Rao, A., Rajesh, G., Ramya, S., & Pai, B. H. (2016). Predictors of cigarette smoking among young adults in Mangalore, India. *Asian Pacific Journal of Cancer Prevention*, 17(1), 45-50. <https://doi.org/10.7314/apjcp.2016.17.1.45>
- Lyons, A., McNeill, A., & Britton, J. (2014). Tobacco imagery on prime time UK television. *Tobacco Control*, 23(3), 257-263. <https://doi.org/10.1136/tobaccocontrol-2012-050650>
- Maretalinia, M., Juliansyah, E., Suyitno, S., Yulianto, A., & Suryani, D. (2021). Association of smoking related to knowledge, attitude, and practice (kap) with tobacco use in Community Health Center Working Area of Sungai Durian, Sintang Regency, West Kalimantan Province. *Buletin Penelitian Sistem Kesehatan*, 24(1), 38-45. <https://doi.org/10.22435/hsr.v24i1.3544>
- Mbatchou Ngahane, B. H., Atangana Ekobo, H., & Kuaban, C. (2015). Prevalence and determinants of cigarette smoking among college students: A cross-sectional study in Douala, Cameroon. *Archives of Public Health*, 73, 47. <https://doi.org/10.1186/s13690-015-0100-1>
- Mbongwe, B., Taper, R., Phaladze, N., Lord, A., & Zetola, N. M. (2017). Predictors of smoking among primary and secondary school students in Botswana. *PLoS One*, 12(4), e0175640. <https://doi.org/10.1371/journal.pone.0175640>
- Ministry of Education Republic of Indonesia. (2015). *The Regulation of Minister of Education of Indonesia Number 64 2015 concerning No Smoking Areas in School Environments stated that school principals, teachers, education staff, students, and other parties are prohibited from smoking*. Jakarta: Ministry of Education Republic of Indonesia.
- Ministry of Finance Republic of Indonesia. (2021). *The Regulation of Minister of Finance Indonesia No 198/PMK 010/2020*. Jakarta: Ministry of Finance Republic of Indonesia.
- Muthukrishnan, A., & Warnakulasuriya, S. (2018). Oral health consequences of smokeless tobacco use.

- Indian Journal of Medical Research*, 148(1), 35-40. https://doi.org/10.4103/ijmr.IJMR_1793_17
- Pengpid, S., & Peltzer, K. (2019). Prevalence and Predictors of Current and Former Tobacco Use among Older Adults in Indonesia. *Asian Pacific Journal of Cancer Prevention*, 20(2), 395-401. <https://doi.org/10.31557/apjcp.2019.20.2.395>
- Prasetyo, B. W., & Adrison, V. (2020). Cigarette prices in a complex cigarette tax system: Empirical evidence from Indonesia. *Tobacco Control*, 29(6), 618-623. <https://doi.org/10.1136/tobaccocontrol-2018-054872>
- Saari, A. J., Kentala, J., & Mattila, K. J. (2014). The smoking habit of a close friend or family member--how deep is the impact? A cross-sectional study. *BMJ Open*, 4(2), e003218. <https://doi.org/10.1136/bmjopen-2013-003218>
- Sinha, D. N., Palipudi, K. M., Rolle, I., Asma, S., & Rinchen, S. (2011). Tobacco use among youth and adults in member countries of South-East Asia region: Review of findings from surveys under the Global Tobacco Surveillance System. *Indian Journal of Public Health*, 55(3), 169-176. <https://doi.org/10.4103/0019-557x.89946>
- Skulberg, K. R., Hamid, S., & Vaktksjold, A. (2019). Smoking among adolescent males at Pulau Weh, Indonesia. *Public Health of Indonesia*, 5(3), 62-70. <https://doi.org/10.36685/phi.v5i3.287>
- Somatunga, L., Sinha, D., Sumanasekera, P., Galapatti, K., Rinchen, S., Kahandaliyanage, A., Mehta, F., & Jayasuriya-Dissanayake, N. (2012). Smokeless tobacco use in Sri Lanka. *Indian Journal of Cancer*, 49(4), 357.
- Statista. (2020). Average retail price of a packet of white cigarettes in Indonesia from 2011 to 2020. <https://www.statista.com/statistics/1182176/indonesia-average-retail-price-of-a-packet-of-cigarettes/>
- Suryati, T., & Tarigan, I. U. (2019). Perilaku konsumsi tembakau pelajar SMP usia 13–15 tahun di Sumatra dan Jawa (Analisis Indonesia–Gyts 2009) [Behavior Tobacco Consumption of Junior High School Student Aged 13–15 Years in Sumatra and Java (Analysis of Indonesia–Gyts 2009)]. *Buletin Penelitian Sistem Kesehatan*, 16(3), 20854.
- Syachroni, S. S., Syarifah, U., Kom, S., & Yulianto, A. (2013). *Riset kesehatan dasar*. Jakarta: Ministry of Health of Indonesia
- Sylvestre, M. P., Hanusaik, N., Berger, D., Dugas, E., Pbert, L., Winickoff, J., & O'Loughlin, J. L. (2018). A tool to identify adolescents at risk of cigarette smoking initiation. *Pediatrics*, 142(5). <https://doi.org/10.1542/peds.2017-3701>
- Thakur, J. S., & Paika, R. (2018). Determinants of smokeless tobacco use in India. *Indian Journal of Medical Research*, 148(1), 41-45. https://doi.org/10.4103/ijmr.IJMR_27_18
- Timberlake, D. S., Nikitin, D., Johnson, N. J., & Altekruse, S. F. (2017). A longitudinal study of smokeless tobacco use and mortality in the United States. *International Journal of Cancer*, 141(2), 264-270. <https://doi.org/10.1002/ijc.30736>
- Tun, N. A., Chittin, T., Agarwal, N., New, M. L., Thuang, Y., & Phyo, P. P. (2017). Tobacco use among young adolescents in Myanmar: Findings from global youth tobacco survey. *Indian J Public Health*, 61(Suppl 1), S54-s59. https://doi.org/10.4103/ijph.IJPH_236_17
- Wibowo, M., Sofiana, L., Ayu, S. M., Gustina, E., & Khoeriyah, I. (2019). Adolescent smoking behaviour determinants in the city of Yogyakarta, Indonesia. *International Journal of Community Medicine and Public Health*, 6(12), 5064-5070. <https://doi.org/10.18203/2394-6040.ijcmph20195445>
- Windiarto, T., Yusuf, A. H., Santoso, A., Nugroho, S., Latifah, S., Solih, R., & Rahmawatiningsih, A. (2018). *Profil anak Indonesia*. Jakarta: Kementerian Pemberdayaan Perempuan dan Perlindungan Anak (KPPA).
- World Health Organization. (2013). *WHO report on the global tobacco epidemic, 2013: Enforcing bans on tobacco advertising, promotion and sponsorship*. Geneva: World Health Organization.
- World Health Organization. (2019). *Tobacco key fact*. Geneva: World Health Organization.
- World Health Organization. (2020). *Global youth tobacco survey (GYTS) Indonesia 2019*. Geneva: World Health Organization.
- Wyss, A. B., Hashibe, M., Lee, Y. A., Chuang, S. C., Muscat, J., Chen, C., Schwartz, S. M., Smith, E., Zhang, Z. F., Morgenstern, H., Wei, Q., Li, G., Kelsey, K. T., McClean, M., Winn, D. M., Schantz, S., Yu, G. P., Gillison, M. L., Zevallos, J. P., . . . Olshan, A. F. (2016). Smokeless Tobacco use and the risk of head and neck cancer: Pooled analysis of US studies in the INHANCE Consortium. *Am J Epidemiol*, 184(10), 703-716. <https://doi.org/10.1093/aje/kww075>
- Zhao, L., Palipudi, K. M., Ramanandraibe, N., & Asma, S. (2016). Cigarette smoking and cigarette marketing exposure among students in selected African countries: Findings from the Global Youth Tobacco Survey. *Preventive Medicine*, 91s, S35-s39. <https://doi.org/10.1016/j.ypmed.2015.12.015>

Cite this article as: Syamsuriansyah., Salman., Rusmitasari, H., Sari, M. H. N., & Maretalinia. (2022). Prevalence and associated factors of Smokeless Tobacco (SLT) use among adolescents in Indonesia: GYTS 2019 data analysis. *Public Health of Indonesia*, 8(1), 9-15. <https://dx.doi.org/10.36685/phi.v8i1.569>